

## Modeling Tree Height, Crown Diameter, Volume and Carbon in Response to Diameter at Breast Height of *Schima wallichii* and *Catanopsis indica*: A Study from Midhills, Nepal

<sup>1</sup>Kabita Koirala, <sup>2</sup>Ram Asheshwar Mandal, <sup>3</sup>Ananda Khadka

<sup>1</sup>Kathmandu Forestry College, Kathmandu, Nepal

<sup>2</sup>School of Environmental Science, Kathmandu, Nepal

<sup>3</sup>Forest Research and Training Centre, Kathmandu, Nepal

**Abstract:** The study regarding correlation between diameter at breast height (DBH) and other variables like height, crown diameter, volume and carbon of the tree species of *Schima wallichii* and *Castanopsis indica* was not done yet. Thus, this research was objectively carried out to assess the correlation between DBH and different parameters particularly height, crown diameter, volume and carbon stock. The study was centered on the mid hill region where natural Schima-Castanopsis forest is abundantly found. Total 360 nested sample plots were established having 12.62 m radius for tree, 5.58 m radius for pole, 2.78 m radius for Sapling, 1.76 m radius for seedling measurement. The DBH and height were measured of these tree species in the plots. Collected data was analyzed using Pearson correlation to show the relation of different characteristics of tree and equations were examined using ANOVA and t-test. The descriptive statistics like mean and standard error (SE) value of DBH was  $28.65 \pm 1.25$  cm, standard deviation was 19.682, minimum value was 5.2 and maximum value was 161.5 of *Castanopsis indica*. Similarly, mean and standard error of DBH was  $22 \pm 0.96$  cm, standard deviation 10.254, minimum value was 5.5 and maximum value was 54.3 of *Schima wallichii*. Moreover, there was positive and strong correlation between DBH and height of *Castanopsis indica* with R square value 0.71 and the equation was  $Y = 7.1075 \ln(X) - 9.4467$  whereas, Y is the height of the tree and X is diameter at breast height. This correlation between DBH and height of *Schima wallichii* also showed positive and strong with R square value 0.81 and the equation was  $Y = 7.9982 \ln(X) - 11.277$ . Same correlation was found between DBH and crown diameter of *Castanopsis indica* having equation  $Y = 3.4335 \ln(X) - 6.8637$  whereas Y is the crown diameter of the tree and X is diameter at breast height. Moreover, similar correlation between DBH and volume was also found. Statistically, ANOVA and t-test showed that there was significant correlation between DBH and tree characteristics like height, crown diameter and volume at 95% confidence level. This research will be useful for the decision maker and scientific community to assess the volume and carbon measuring single variable i.e. only DBH.

**Key words:** DBH • Height • Basal area • Volume • Crown diameter

### INTRODUCTION

*Schima wallichii* and *Castanopsis indica* are hilly plant species generally found in Bangladesh, Bhutan, southwestern China, Myanmar, Thailand, Laos, Vietnam, northern India and Nepal [1]. Both species are medium fast growing. *Schima wallichii* is an evergreen tree belonging to the tea family *Theaceae* while *Castanopsis indica* is belonging to *Fagaceae* family. These species

are used mainly for timber to construct the house in hilly areas in these countries. The leaves of these species are used for fodder and nuts are edible. The mid hill is very suitable habitat for this species. These species generally distributed in eastern central hilly region of Nepal [2].

Several types of models are essential to understand tree characteristics [3, 4, 5]. These models are differed according to species and their use [6,7]. The timber is most valuable forest product which is traded in volume

unit [8, 9, 10]. Similarly, another important product is biomass and carbon [11]. Carbon sequestration is vital role of tree which is also known as carbon credit [12, 13, 14]. This carbon sequestration helps us to understand the phenomenon of climate change [8, 15, 16]. The volume and carbon assessment of felled trees is easy and convenient job but it is very difficult to assess these parameters of standing trees [17]. Thus, reliable equations are needed. There are very limited equations to assess the volume and carbon in Nepal.

The diameter is most important variable in the forest science. Almost all parameters correlate with the diameter at breast. There is strong correlation between diameter and other variables like height, volume and carbon. However, these correlations are varied according to species, climatic condition and geographical areas. The correlations of these variables of *Schima wallichii* and *Castanopsis indica* are limitedly explored in Nepal. The assessment of correlation between the variables is important to show the height, volume and carbon of the tree with respect to diameter at breast height. These equations will help to evaluate volume and carbon stock only measuring the diameter at breast. Thus, such types of research are essential. Therefore, this study was carried out to assess the correlation between DBH and different parameters like height, volume and carbon stock.

## MATERIALS AND METHODS

**Study Area:** The study was centered on the mid hill region where natural *Schima-Castanopsis* forest is dominantly found. Mid Hills, lie north of Churia along the southern flanks of the Himalayas [18]. The region occupies 4, 306, 230 ha (29.2%) of the total land area of the country and covers parts of 55 of the nation's districts. The specific plots for the study were determined after the preliminary visit and study (Figure 1).

**Sampling Design:** Permanent sample plots (PSP) were used for inventory works. The sample plots were chosen in such a way that environmental variability can be incorporated into the growth of species. Plot sizes were chosen according to the guideline of Forest Research and Training Centre (FRTC) of Nepal and measurement was done.

**Data Collection and Analysis:** Total 360 nested sample plots were established to collect the data. The plots were established having 12.62 m radius for tree, 5.58 m radius for pole, 2.78 m radius for Sapling, 1.76 m radius for seedling measurement. The DBH and height were measured of these tree species in the forest.

The following formulae were used for the quantitative analysis.




Fig. 1: Mid-hill (Middle Mountain) Region of Nepal

**Volume Calculation:**

$$Volume (V) = \left(\frac{\pi d^2}{4} * Height * from factor\right) / 10000$$

whereas, V = Volume in m<sup>3</sup> and d = diameter in cm

**Above Ground Tree Biomass (AGTB) Calculation:** The equation developed by Chave *et al.* [19] was used to calculate above ground tree biomass,

$$AGTB = 0.0509 * \rho * d^2 * H$$

where,

AGTB = above ground tree biomass (kg),

$\rho$  = wood density of the species (gm/cm<sup>3</sup>)

d = diameter (cm) and H = tree height (m)

**Carbon Calculation:** Again, the total biomass stock was converted into carbon stocks using conversion factor 0.47 [20].

$$Carbon (C) = 0.47 * total\ biomass$$

Collected data was analyzed using Pearson correlation to show the relation of different characteristics of tree and these equations were examined using ANOVA and t-test.

**RESULTS**

**Descriptive Statistics of Different Parameters of *Castanopsis indica* and *Schima wallichii*:** The descriptive statistics includes mean, standard error, standard deviation, minimum and maximum value of diameter, carbon, height and crown diameter of *Castanopsis indica*. The mean and standard error (SE) value of DBH was 28.65±1.25cm, standard error was 1.25, standard deviation 19.682, minimum value was 5.2 and maximum value was 161.5. Similar values were estimated for height, crown diameter, volume and carbon of *Schima wallichii*. The mean and standard error value of DBH was 22±0.96 cm, standard error was 0.96, standard deviation 10.254, minimum value was 5.5 and maximum value was 54.3. Similar values were estimated for height, crown diameter, volume and carbon.

**Correlation Between DBH and Height of *Castanopsis indica* as well as *Schima wallichii*:** The result showed that there was positive and strong correlation

between DBH and height of *Castanopsis indica*. The equation was  $Y=7.1075\ln(X)-9.4467$  whereas, Y is the height of the tree and X is diameter at breast height. The estimated R-square value was 0.7117 (Figure 2). Similar correlation was found between DBH and height of *Schima wallichii*. The equation was  $Y=7.9982\ln(X)-11.277$  whereas, Y is the height of the tree and X is diameter at breast height. The estimated R-square value was 0.811 (Figure 3).

Statistically, ANOVA showed that there was significant relation between DBH and height. In addition, t-test also showed that intercept and x-variable of the equation was significant since P-value was less than 0.05 in both test.

**Correlation Between DBH and Crown Diameter of *Castanopsis indica* as well as *Schima wallichii*:** The correlation between DBH and crown diameter of *Castanopsis indica* was also strong and positive. The derived equation was  $Y=3.4335\ln(X)-6.8637$  whereas Y is the crown diameter of the tree and X is diameter at breast height. The estimated R-square value was 0.8241 (Fig. 4). This result was also same to show correlation between DBH and crown diameter of *Schima wallichii* but the equation was different, i.e.  $Y=11.803\ln(X)-25.663$  whereas, Y is the crown diameter of tree and X is diameter at breast height. The estimated R-square value was 0.8462 (Figure 5).

The equation and its intercept and x-variable were significant at 95% confidence level, as the p-value of ANOVA and t-test showed less than 0.05.

**Correlation Between DBH and Volume of *Castanopsis indica* and *Schima wallichii*:** It was positive and strong correlation between DBH and volume of *Castanopsis indica*. The equation was  $Y=0.1304\ln(X)-2.6737$  whereas, Y is the volume of *Castanopsis indica* and X is the diameter at breast height. The estimated R-square value was 0.7614 (Figure 6).

Similar correlation was found between DBH and volume of *Schima wallichii*. The equation was  $Y = 0.0424\ln(X)-0.566$  whereas, Y is the volume of *Schima wallichii* and X is the diameter at breast height. The estimated R square value was 0.8419 (Figure 7). ANOVA and t-test showed that the equation showing correlation between DBH and volume of *Castanopsis indica* and *Schima wallichii* was significant, the intercept as well as x-variable was also significant since p-values of these tests was less than 0.05.

Table 1: Descriptive statistics of characteristics of *Castanopsis indica* and *Schima wallichii*

Statistics	Mean±SE	Standard deviation	Minimum value	Maximum value
<i>Castanopsis indica</i>				
DBH	28.65±1.25	19.682	5.2	161.5
Height	13.13±0.32	5.018	3.9	32.8
Crown diameter	4±0.143	2.252	1.4	15
Volume	1±0.187	2.94	0	32.96
Carbon	1464.83±259.56	4056.74	9.45	45472.1
<i>Schima wallichii</i>				
DBH cm	22±0.96	10.25	5.5	54.3
Height m	12±0.40	4.29	3	25
Crown diameter	9.5±0.58	6.19	1	20.5
Volume	0.3±0.04	0.47	0	2.5
Carbon	509±60	653.15	6.45	3576.57




Fig. 2: Correlation between DBH&ht of *Castanopsis indica*




Fig. 3: Correlation between DBH&ht of *Castanopsis indica*

**Correlation Between DBH and Carbon Stock of *Castanopsis indica* and *Schima wallichii*:** The correlation between DBH and carbon stock of *Castanopsis indica* was also positive and strong. The equation was  $Y=179.85\ln(X)-3688.6$  whereas, Y is the carbon stock of *Castanopsis indica* and X is the diameter at breast height.

The estimated R-square value was 0.7614 (Figure 8). Similar correlation was found between DBH and carbon stock of *Schima wallichii*. The equation was  $Y=58.445\ln(X)-780.91$  whereas, Y is the carbon stock of *Schima wallichii* and X is the diameter at breast height. The estimated R-square value was 0.8419.




Fig. 4: Correlation between DBH and crown diameter of *Castonopsis indica*




Fig. 5: Correlation between DBH and crown diameter of *Schima wallichii*




Fig. 6: Correlation between DBH and volume of *Castonopsis indica*




Fig. 7: Correlation between DBH and volume of *Schima wallichii*




Fig. 8: Correlation between DBH and carbon of *Castonopsis indica*




Fig. 9: Correlation between DBH and carbon of *Schima wallichii*

The p-value of ANOVA and t-test analysis showed that the carbon estimation equations of *Castonopsis indica* and *Schima wallichii* was significant, the intercept as well as x-variable was also significant since p-values of these tests was less than 0.05.

### DISCUSSION

There was positive and strong correlation between DBH and height [21, 22, 23] of *Castonopsis indica* and *Schima wallichii*. This research also showed an excellent relationship between DBH and tree height. Several authors support this concept. Chaturvedi and Khan, [24], Huang *et al.* [25] and many authors showed this relation. The findings of this research showed that, as the diameter

increases the height also increase but at a certain point height growth is constant. This concept is also supported by West and West [26].

The most common use variables are diameter at breast height, height, form factors and site quality [27, 28, 29] to determine the volume, crown diameter and carbon. This model is based on diameter at breast height. The height, crown diameter, volume and carbon models were developed with respect to DBH. Similar variables are used to determine the height, crown diameter, volume and carbon [30, 31, 32].

The correlation between tree characteristics like height, volume and carbon stock and DBH was affected due to several factors [33, 34, 35]. Overall, fire was the most prominent cause in all community forests followed

by litter. In fact, intentional fire was general cause of the forests carbon loss. The report of DFRS [18] also showed that grazing as the most disturbing factor in carbon stock. The report published on drivers of deforestation and forest degradation showed illegal logging, grazing and fire were the main drivers of carbon loss [36].

### CONCLUSION AND RECOMMENDATION

The correlation between diameter and other parameter like height, volume, crown diameter and carbon stock was strong and positive. The ANOVA and t-test showed that the correlation between diameter and other parameter was significant. These equations can be used to assess the height, volume, crown diameter and carbon stock only measuring single parameter i.e. diameter at breast height. Similarly, these equations will be useful to derive other equation at local level as well. The research should be carried out to show the relation between DBH and other parameters as well as how these relations response to factors likes aspect, altitude and soil.

### ACKNOWLEDGEMENT

We would like to acknowledge for technical support provided by Forest Research and Training Centre and Kathmandu Forestry College, Nepal.

### REFERENCES

1. Bhandari, A.R., U.R. Khadka and K.R. Kanel, 2018. Ecosystem services in the mid-hill forest of western Nepal: A case of Panchase protected forest. *Journal of Institute of Science and Technology*, 23(1), 10-17.
2. Gautam, C.M. and T. Watanabe, 2005. Composition, distribution and diversity of tree species under different management systems in the hill forests of Bharse Village, Gulmi District, Western Nepal. *Himalayan Journal of Sciences*, 3(5), 67-74.
3. Kempes, C.P., G.B. West, K. Crowell and M. Girvan, 2011. Predicting maximum tree heights and other traits from allometric scaling and resource limitations. *PLoS One*, 6(6): e20551.
4. Mandal, R.A., I.C. Dutta, P.K. Jha and S. Karmacharya, 2013. Relationship between Carbon Stock and Plant Biodiversity in Collaborative Forests in Terai, Nepal [Research Article]. *ISRN Botany; Hindawi*. <https://doi.org/10.1155/2013/625767>.
5. Temesgen, H., D. Affleck, K. Poudel, A. Gray and J. Sessions, 2015. A review of the challenges and opportunities in estimating above ground forest biomass using tree-level models. *Scandinavian Journal of Forest Research*, 30(4): 326-335.
6. Condit, R., 2008. Methods for estimating aboveground biomass of forest and replacement vegetation in the tropics. *Center for Tropical Forest Science Research Manual*, 73: 1-73.
7. Köhl, M., R. Lasco, M. Cifuentes, O. Jonsson, K.T. Korhonen, P. Mundhenk, J. De Jesus Navar and G. Stinson, 2015. Changes in forest production, biomass and carbon: Results from the 2015 UN FAO Global Forest Resource Assessment. *Forest Ecology and Management*, 352: 21-34. <https://doi.org/10.1016/j.foreco.2015.05.036>.
8. Seidl, R., W. Rammer, D. Jäger, W.S. Currie and M.J. Lexer, 2007. Assessing trade-offs between carbon sequestration and timber production within a framework of multi-purpose forestry in Austria. *Forest Ecology and Management*, 248(1-2): 64-79.
9. Kurz, W.A., C.C. Dymond, T.M. White, G. Stinson, C.H. Shaw, G.J. Rampley and M.J. Apps, 2009. CBM-CFS3: a model of carbon-dynamics in forestry and land-use change implementing IPCC standards. *Ecological Modelling*, 220(4): 480-504.
10. Powell, S.L., W.B. Cohen, S.P. Healey, R.E. Kennedy, G.G. Moisen, K.B. Pierce and J.L. Ohmann, 2010. Quantification of live aboveground forest biomass dynamics with Landsat time-series and field inventory data: A comparison of empirical modeling approaches. *Remote Sensing of Environment*, 114(5): 1053-1068. <https://doi.org/10.1016/j.rse.2009.12.018>.
11. Hunt, C.A.G., 2009. *Carbon Sinks and Climate Change: Forests in the Fight Against Global Warming*. Edward Elgar Publishing.
12. Banskota, K., B. Karky and M. Skutsch, 2007. Reducing carbon emissions through community-managed forests in the Himalaya. *Reducing Carbon Emissions through Community-Managed Forests in the Himalaya*. <https://www.cabdirect.org/cabdirect/abstract/20083078706>.
13. Gonzalez, P., J.J. Battles, B.M. Collins, T. Robards, and D.S. Saah, 2015. Aboveground live carbon stock changes of California wildland ecosystems, 2001-2010. *Forest Ecology and Management*, 348, 68-77. <https://doi.org/10.1016/j.foreco.2015.03.040>.
14. Mandal, R.A.I.C., Dutta, P.K. Jha and S.B. Karmacharya, 2016. Carbon sequestration potential in community and collaborative forests in Terai, Nepal. [/paper/Carbon-sequestration-potential-in-community-and-in-andalDutta/082675ee04036398ff8518117b712412324b3515](https://doi.org/10.1155/2016/20162016).
15. Gautam, K.R., 2002. Carbon sequestration in agroforestry and annual cropping system in inner Terai, Central Nepal. <https://agris.fao.org/agrissearch/search.do?recordID=NO2002197591>.

16. Baral, S.K., R. Malla and S. Ranabhat, 2009. Above-ground carbon stock assessment in different forest types of Nepal. *BankoJanakari*, 19(2): 10-14. <https://doi.org/10.3126/banko.v19i2.2979>.
17. Vallet, P., J.F. Dhôte, G. Le Moguédec, M. Ravart and G. Pignard, 2006. Development of total aboveground volume equations for seven important forest tree species in France. *Forest Ecology and Management*, 229(1-3): 98-110.
18. DFRS, 2015. States of Nepal's Forests, Department of Forest Research and Survey and Forest Resource Assessment (DFRS and FRA) Nepal, pp: 34-45.
19. Chave, J., C. Andalo, S. Brown, M.A. Cairns, J.Q. Chambers, D. Eamus, H. Fölster, F. Fromard, N. Higuchi, T. Kira, J.P. Lescure, B.W. Nelson, H. Ogawa, H. Puig, B. Riéra and T. Yamakura, 2005. Tree allometry and improved estimation of carbon stocks and balance in tropical forests. *Oecologia*, 145(1):87-99. <https://doi.org/10.1007/s00442-005-0100-x>.
20. Gairola, S., C.M. Sharma, S.K. Ghildiyal and S. Suyal, 2011. Live tree biomass and carbon variation along an altitudinal gradient in moist temperate valley slopes of the Garhwal Himalaya (India). *Current Science*, 1862-1870.
21. Kohyama, T., E. Suzuki, T. Partomihardjo, T. Yamada and T. Kubo, 2003. Tree species differentiation in growth, recruitment and allometry in relation to maximum height in a Bornean mixed dipterocarp forest. *Journal of Ecology*, 91(5): 797-806.
22. Buba, T., 2013. Relationships between stem diameter at breast height (DBH), tree height, crown length and crown ratio of *Vitellariaparadoxa* CF Gaertn in the Nigerian Guinea Savanna. *African Journal of Biotechnology*, 12(22).
23. Shrestha, S., B.S. Karky, A. Gurung, R. Bista and O.R. Vetaas, 2013. Assessment of Carbon Balance in Community Forests in Dolakha, Nepal. *Small-Scale Forestry*, 12(4): 507-517. <https://doi.org/10.1007/s11842-012-9226-y>.
24. Khanna, L.S. and A.N. Chaturbedi, 1982. *Forest Mensuration*. International Book Distributer, Deharadun, India.
25. Huang, X., B. Ziniti, N. Torbick and M.J. Ducey, 2018. Assessment of forest above ground biomass estimation using multi-temporal C-band sentinel-1 and polarimetric L-band PALSAR-2 data. *Remote Sensing*, 10(9): 1424.
26. Mehtätalo, L., S. de-Miguel and T.G. Gregoire, 2015. Modeling height-diameter curves for prediction. *Canadian Journal of Forest Research*, 45(7): 826-837.
27. King, D.A., S.J. Davies, S. Tanand N.S.M. Noor, 2006. The role of wood density and stem support costs in the growth and mortality of tropical trees. *Journal of Ecology*, 94(3): 670-680.
28. Chapagain, T.R. and R.P. Sharma, 2021. Modeling form factors for sal (*Shorea robusta* Gaertn.) using tree and stand measures, and random effects. *Forest Ecology and Management*, 482, 118807.
29. Mandal, R.A., B.K.V. Yadav, K.K. Yadav, I.C. Dutta and S.M. Haque, 2013. Development of allometric equation for biomass estimation of eucalyptus *camaldulensis*: a study from Sagarnath Forest, Nepal. *Int. J. Biodiv. Ecosyst.*, 1: 1-7.
30. Pavan, M.N., K.S. Sunil and A. Ghosh, 2022. (forthcoming). Analysis of *E. regnans* Form Characteristics. IEEE 11<sup>th</sup> International Conference on Communication Systems & Network Technologies (CSNT) (pp. 329-334). IEEE.
31. Egeta, D., M. Negash, M. Alebachew, S. Mulgeta and T. Lemi, 2022 (forthcoming). Species specific allometric equations, biomass expansion factor and wood density of native tree species in dry afro-montane forest of Ethiopia.
32. Mäkelä, Annikki. 2002. Derivation of stem taper from the pipe theory in a carbon balance framework. *Tree Physiology*, 22(13):891-905.
33. Malhi, Y. and J. Grace, 2000. Tropical forests and atmospheric carbon dioxide. *Trends in Ecology & Evolution*, 15(8): 332-337. [https://doi.org/10.1016/S01695347\(00\)01906-6](https://doi.org/10.1016/S01695347(00)01906-6).
34. Luysaert, S., E.D. Schulze, A. Börner, A. Knohl, D. Hessenmöller, B.E. Law, P. Ciais and J. Grace, 2008. Old-growth forests as global carbon sinks. *Nature*, 455(7210): 213-215. <https://doi.org/10.1038/nature07276>.
35. Domke, G.M., C.W. Woodall, J.E. Smith, J.A. Westfall and R.E. McRoberts, 2012. Consequences of alternative tree-level biomass estimation procedures on US forest carbon stock estimates. *Forest Ecology and Management*, 270: 108-116.
36. Geist, J. and F. Lambin, 2001. What Drives Tropical Deforestation? A meta-analysis of proximate and underlying causes of deforestation based on subnational case study evidence, LUCR Report Series 4, CIACO, Louvain-la-Neuve, pp: 138-142.